

$t\bar{t}$ production and decay at NNLO QCD

Rene Poncelet¹

Institut für Theoretische Teilchenphysik und Kosmologie

2016-08-30



¹in cooperation with M. Czakon

Introduction

The amplitudes

Factorization of decays

$t\bar{t}$ production amplitudes


Structure of born-like amplitudes

Renormalization and finite remainder

Summary

Introduction

Why is the top-quark (still) interesting?




Recent developments

Stable onshell tops and spin summed:

- Total inclusive cross sections @ NNLO+NNLL accuracy [Czakon, Fiedler, Mitov '13]
- Fully differential distributions @ NNLO

Unstable tops + spin correlations:

- Off-shell effects with decays @ NLO + matched parton-shower




Recent developments

Stable onshell tops and spin summed:

- Total inclusive cross sections @ NNLO+NNLL accuracy [Czakon, Fiedler, Mitov '13]
- Fully differential distributions @ NNLO

Unstable tops + spin correlations:

- Off-shell effects with decays @ NLO + matched parton-shower




Recent developments

Stable onshell tops and spin summed:

- Total inclusive cross sections @ NNLO+NNLL accuracy [Czakon, Fiedler, Mitov '13]
- Fully differential distributions @ NNLO

Unstable tops + spin correlations:

- Off-shell effects with decays @ NLO + matched parton-shower




The amplitudes

Feynman diagrams

process in mind: $pp \rightarrow t\bar{t}$


production:



Feynman diagrams


process in mind: $t \rightarrow b + 2f$

production:



decay:


+



Feynman diagrams


process in mind: $pp \rightarrow t\bar{t} \rightarrow b\bar{b} + 4f$

production:




+

decay:



=


combined:



Feynman diagrams


process in mind: $pp \rightarrow t\bar{t} \rightarrow b\bar{b} + 4f$

production:




+

decay:




=

combined:




Feynman diagrams

$$pp \rightarrow b\bar{b} + 4f$$



Feynman diagrams

$$pp \rightarrow b\bar{b} + 4f$$




Simplification I

restrict to onshell tops

- Narrow-Width-Approximation
- Factorization of production and decay
- Much simpler production amplitudes
- Neglecting continuum production

Feynman diagrams

$$pp \rightarrow t\bar{t} \rightarrow b\bar{b} + 4f$$




Simplification I

restrict to onshell tops


- Narrow-Width-Approximation
- Factorization of production and decay
- Much simpler production amplitudes
- Neglecting continuum production

Factorization of decays


Narrow-Width-Approximation



Narrow-Width-Approximation



Narrow-Width-Approximation



- enters matrix element as:

$$\sim \frac{1}{(p^2 - m^2)^2 + m^2\Gamma^2}$$


- For crosssections: Integration over phase-space
- + limit $\Gamma/m \rightarrow 0$:

$$\frac{1}{(p^2 - m^2)^2 + m^2\Gamma^2} \rightarrow \frac{2\pi}{2m\Gamma} \delta(p^2 - m^2)$$

- On amplitude level:

$$\mathcal{M} = \mathcal{M}_{\text{NWA}} + \mathcal{O}\left(\frac{\Gamma}{m}\right)$$

Amplitude factorization



$$\mathcal{M} = \left(\tilde{A}(t \rightarrow bl^+ \nu) \frac{i(\not{p}_t + m)}{p_t^2 - m^2 + im\Gamma_t} \right) \cdot \tilde{A}(pp \rightarrow \bar{t}t) \cdot \left(\frac{i(-\not{p}_{\bar{t}} + m)}{p_{\bar{t}}^2 - m^2 + im\Gamma_t} \tilde{A}(\bar{t} \rightarrow \bar{b}l^- \bar{\nu}) \right)$$

Decay spinors

Narrow-Width-Approximation:

$$\frac{i(-\not{p}_{\bar{t}} + m)}{p_{\bar{t}}^2 - m^2 + im\Gamma_t} \tilde{A}(\bar{t} \rightarrow \bar{b}l^{-}\bar{\nu}) \rightarrow \frac{i(-\not{p}_{\bar{t}} + m)}{\sqrt{2m\Gamma_t}} \tilde{A}(\bar{t} \rightarrow \bar{b}l^{-}\bar{\nu})$$


Decay spinors

$$\bar{U}(p_t) = \tilde{A}(t \rightarrow bl^+\nu) \frac{i(\not{p}_t + m)}{\sqrt{2m\Gamma_t}}$$
$$V(p_{\bar{t}}) = \frac{i(-\not{p}_{\bar{t}} + m)}{\sqrt{2m\Gamma_t}} \tilde{A}(\bar{t} \rightarrow \bar{b}l^{-}\bar{\nu})$$


Amplitude:

$$\mathcal{M} = \bar{U}(p_t) \tilde{A}(pp \rightarrow \bar{t}t) V(p_{\bar{t}}) + \mathcal{O}\left(\frac{\Gamma_t}{m}\right)$$

QCD corrections to decay




QCD corrections to decay



QCD corrections to decay

Simplification II

restrict to leptonic top decays



- Vertex corrections (for massless final state):

$$\Gamma^\mu = \frac{g}{\sqrt{2}} \left\{ \gamma^\mu [F_{1L}P_L + F_{1R}P_R] + \frac{i\sigma^{\mu\nu}q_\nu}{2m_t} [F_{2L}P_R + F_{2R}P_L] \right\}$$

- times W propagator and decay vertex

$$\bar{u}(p_\nu) \frac{ig_W}{\sqrt{2}} \gamma^\nu \frac{(1 - \gamma_5)}{2} v(p_{l^+}) \cdot \frac{-i(g_{\nu\mu} - \frac{q_\nu q_\mu}{q^2})}{q^2 - m_W^2 + i\Gamma_W m_W} \bar{u}(p_b) i\Gamma^\mu$$

Contributions to amplitude

Decay \ Production	LO	NLO	NNLO
LO	Green	Green	Green
NLO	Green	Green	Red
NNLO	Green	Red	Red

Contributions to amplitude

Production \ Decay	LO	NLO	NNLO
LO			
NLO			
NNLO			

$t\bar{t}$ production amplitudes

$t\bar{t}$ production amplitudes

Goal: Evaluate $\bar{U}(p_{\bar{t}})\tilde{A}(pp \rightarrow \bar{t}t)V(p_{\bar{t}})$




$t\bar{t}$ production amplitudes

Goal: Evaluate $\bar{U}(p_t)\tilde{A}(pp \rightarrow \bar{t}t)V(p_{\bar{t}})$

$t\bar{t}$ production amplitudes

Goal: Evaluate $\bar{U}(p_t)\tilde{A}(pp \rightarrow \bar{t}t)V(p_{\bar{t}})$

Contributions to $\tilde{A}(pp \rightarrow \bar{t}t)$:

		gg qq
LO		3 1
NLO		33 18
NNLO		726 190

Structure of born-like amplitudes

Lorentz-structures – Gluon channel

Lorentz-decomposition: $\mathcal{A} = \epsilon_{1\mu}(p_1)\epsilon_{2\nu}(p_2)A^{\mu\nu}$

$A^{\mu\nu}$ is a rank-2 Lorentz tensor

Using:

- momentum conservation
- transversality
- equation of motion

leads to 10 independent structures in $d = 4 - 2\epsilon$ dimensions

$$A^{\mu\nu} = \sum_{j=1}^{10} A_j T_j^{\mu\nu}$$

$$T_1^{\mu\nu} = m\bar{u}_3\gamma^\mu\gamma^\nu v_4$$

$$T_2^{\mu\nu} = \bar{u}_3\gamma^\mu v_4 p_3^\nu$$

$$T_3^{\mu\nu} = \bar{u}_3\gamma^\nu v_4 p_3^\mu$$

$$T_4^{\mu\nu} = m\bar{u}_3 v_4 g^{\mu\nu}$$

$$T_5^{\mu\nu} = m^{-1}\bar{u}_3 v_4 p_3^\mu p_3^\nu$$

$$T_6^{\mu\nu} = m^{-2}\bar{u}_3 \not{p}_1 v_4 p_3^\mu p_3^\nu$$


$$T_7^{\mu\nu} = \bar{u}_3 \not{p}_1 v_4 g^{\mu\nu}$$

$$T_8^{\mu\nu} = m^{-1}\bar{u}_3 \not{p}_1 \gamma^\nu v_4 p_3^\mu$$

$$T_9^{\mu\nu} = m^{-1}\bar{u}_3 \not{p}_1 \gamma^\mu v_4 p_3^\nu$$

$$T_{10}^{\mu\nu} = \bar{u}_3 \not{p}_1 \gamma^\mu \gamma^\nu v_4$$

Lorentz-structures – Quark channel



- two disconnected fermion lines
- connection by gluons+loops
- parity conservation \rightarrow no γ_5

$$\tilde{A} = \sum_{i=1}^7 A_i T_i$$

$$\text{with } T_j \sim \bar{v}_2 \Gamma_j u_1 \bar{u}_3 \Gamma'_j v_4$$

$$T_1 = m^{-1} \bar{v}_2 \not{p}_3 u_1 \bar{u}_3 v_4$$

$$T_2 = m^{-2} \bar{v}_2 \not{p}_3 u_1 \bar{u}_3 \not{p}_1 v_4$$

$$T_3 = \bar{v}_2 \gamma^\mu u_1 \bar{u}_3 \gamma_\mu v_4$$

$$T_4 = \bar{v}_2 \gamma^\mu u_1 \bar{u}_3 \not{p}_1 \gamma_\mu v_4$$

$$T_5 = m^{-1} \bar{v}_2 \not{p}_3 \gamma^\mu \gamma^\nu u_1 \bar{u}_3 \gamma_\mu \gamma_\nu v_4$$

$$T_6 = m^{-2} \bar{v}_2 \not{p}_3 \gamma^\mu \gamma^\nu u_1 \bar{u}_3 \not{p}_1 \gamma_\mu \gamma_\nu v_4$$

$$T_7 = \bar{v}_2 \gamma^\mu \gamma^\nu \gamma^\rho u_1 \bar{u}_3 \not{p}_1 \gamma_\mu \gamma_\nu \gamma_\rho v_4$$

color decomposition: $\mathcal{A} = \sum_{ij} c_{ij} C_i A_j$

Gluon channel

- Gluons: a, b adjoint rep.
- Quarks: c, d fundamental rep.

$$C_1 = (T^a T^b)_{cd}$$

$$C_2 = (T^b T^a)_{cd}$$

$$C_3 = \text{Tr} \{ T^a T^b \} \delta_{cd}$$

Quark channel

- Quarks: a, b fundamental rep.
- Quarks: c, d fundamental rep.

$$C_1 = \delta_{ac} \delta_{bd}$$

$$C_2 = \delta_{ab} \delta_{cd}$$

Projection

Idea: Construct projectors: $P_j = \sum_l B_{jl}(T_l)^\dagger$

Extracting the B_{jl} :

$$\sum_{spin/pol/col} P_j \mathcal{A} \stackrel{!}{=} A_j$$

leads to system of equations

$$\sum_{l,k} B_{jl} A_k \sum_{spin/pol/col} (T_l)^\dagger T_k = A_j$$

Inversion \rightarrow coefficients B_{jl}

Projection

Idea: Construct projectors: $P_j = \sum_l B_{jl}(T_l)^\dagger$

Extracting the B_{jl} :

$$\sum_{spin/pol/col} P_j \mathcal{A} \stackrel{!}{=} A_j$$

leads to system of equations

$$\sum_{l,k} B_{jl} A_k \sum_{spin/pol/col} (T_l)^\dagger T_k = A_j$$

Inversion \rightarrow coefficients B_{jl}

Example: Lorentz-structures gluons

$$\sum_{pol, spin} P_j^{\mu_1 \nu_1} \left[\epsilon_{1\mu_1}^* \epsilon_{2\nu_1}^* \epsilon_{1\mu_2} \epsilon_{2\nu_2} \right] A^{\mu_2 \nu_2} =$$


$$\text{Tr} \left[(\not{p}_4 - m) P_j^{\mu_1 \nu_1} (\not{p}_3 + m) \cdot \right.$$

$$\left. \left(-g_{\mu_1 \mu_2} + \frac{p_{2\mu_1} p_{1\mu_2} + p_{2\mu_2} p_{1\mu_1}}{p_1 \cdot p_2} \right) \cdot \right.$$

$$\left. \left(-g_{\nu_1 \nu_2} + \frac{p_{1\nu_1} p_{2\nu_2} + p_{1\nu_2} p_{2\nu_1}}{p_1 \cdot p_2} \right) A^{\mu_2 \nu_2} \right]$$

Summary


$$\bar{U}(p_t)\tilde{A}(pp \rightarrow \bar{t}t)V(p_{\bar{t}})$$




Summary

$$\bar{U}(p_t) \tilde{A}(pp \rightarrow \bar{t}t) V(p_{\bar{t}})$$

- NWA \rightarrow decay spinors



Summary




$$\bar{U}(p_t)\tilde{A}(pp \rightarrow \bar{t}t)V(p_{\bar{t}})$$

- NWA \rightarrow decay spinors
- Lorentz- and color-decomposition

$$\tilde{A} = \sum_{ij} c_{ij} C_i \tilde{A}_j$$

Summary



$$\bar{U}(p_t)\tilde{A}(pp \rightarrow \bar{t}t)V(p_{\bar{t}})$$

- NWA \rightarrow decay spinors
- Lorentz- and color-decomposition

$$\tilde{A} = \sum_{ij} c_{ij} C_i \tilde{A}_j$$

- Evaluation of the coefficients
 - functions of $ms = m^2/s, x = t/s, \epsilon$
 - and scalar 2-loop integrals
 - \rightarrow reuse of numerical integrals from spin summed calculation

Renormalization and finite remainder

UV renormalization and decoupling

$$|M_{g,q}(\alpha_S^0, m^0, \epsilon)\rangle = 4\pi\alpha_S^0 \left[|M_{g,q}^{(0)}(m^0, \epsilon)\rangle + \left(\frac{\alpha_S^0}{2\pi}\right) |M_{g,q}^{(1)}(m^0, \epsilon)\rangle + \left(\frac{\alpha_S^0}{2\pi}\right)^2 |M_{g,q}^{(2)}(m^0, \epsilon)\rangle \right]$$

UV-renormalized amplitude:

$$|\mathcal{M}_{g,q}^R(\alpha_S^{(n_f)}(\mu), m, \mu, \epsilon)\rangle = \left(\frac{\mu^2 e^{\gamma_E}}{4\pi}\right)^{-2\epsilon} Z_{g,q} Z_Q |M_{g,q}(\alpha_S^0, m^0, \epsilon)\rangle$$

- Z_g, Z_q, Z_Q : onshell renormalization constants
- $m^0 = Z_m m$
- $\alpha_S^0 = \left(\frac{e^{\gamma_E}}{4\pi}\right)^\epsilon \mu^{2\epsilon} Z_{\alpha_S}^{(n_f)} \alpha_S^{(n_f)}(\mu)$
 $\hat{=} \overline{MS}$ -scheme with n_f flavours

Decoupling

- $n_f = n_l + n_h$ is not feasible
- decouple the running of α_S from the n_h quarks
- $\alpha_S^{(n_f)} = \zeta_{\alpha_S} \alpha_S^{(n_l)}$

IR divergences and the finite remainder function

$$\left| \mathcal{M}_{g,q}^{fin} \left(\alpha_S^{(n_f)}(\mu), m, \mu, \epsilon \right) \right\rangle = \mathbf{Z}_{\mathcal{M}_{g,q}}^{-1}(\{p\}, m, \mu, \epsilon) \left| \mathcal{M}_{g,q}^R \left(\alpha_S^{(n_f)}(\mu), m, \mu, \epsilon \right) \right\rangle$$

- Amplitude still contains infrared divergences
- Usually: Cancellation with divergences from real-emission contributions
- Complete factorization of IR structure $\rightarrow \mathbf{Z}$ operator
- \rightarrow finite remainder

- Z can be calculated by its anomalous dimension equation

$$\frac{d}{d \ln \mu} \mathbf{Z} = -\Gamma \mathbf{Z}$$

- depends on kinematics
- matrix in color space

Summary of progress

Finished

- Projection LO, NLO, NNLO amplitudes
- Finite remainder of all coefficients
- Implementation of Color and Spin Structures in STRIPPER
- Implementation of decay spinors in STRIPPER
- Implementation of LO, NLO coefficients in STRIPPER


Outlook

- Implementation of NNLO finite remainder
- Real-Virtual and Real-Real contributions
- Implementation of decay phase-space and handling of decay products in STRIPPER
- Merging with the results from Mitov et al.
→ QCD corrections to decay

Thank you for your attention.


Backup

Recent results: Measurements



Top quark results from the LHC | Nuno Castro | QCD@LHC 2016

Recent results: Measurements



STRIPPER – SecToR Improved Phase sPacE for real Radiation

The subtraction scheme

- Method of evaluate the double-real emission radiation contribution to NNLO processes
- Decomposition of the phase-space to factorize the singular limits of the amplitude
- Suitable parameterizations to derive (integrated) subtraction terms

The NNLO event generator


- fully differential event generation
- several scales simultaneously
- different pdfs simultaneously
- stable tops
- predecided binned distributions
- fixed top mass=173.3

Why precision predictions for top-quark pair production? → Applications

- top quark mass and production cross-sections
→ precision test of SM
- measurements of α_S
- Constrains on PDFs
- SM phase diagram
- BSM searches in production and decay


Why precision predictions for top-quark pair production? → Applications

- top quark mass and production cross-sections
→ precision test of SM
- **measurements of α_S**
- Constrains on PDFs
- SM phase diagram
- BSM searches in production and decay





Why precision predictions for top-quark pair production? → Applications

- top quark mass and production cross-sections
→ precision test of SM
- measurements of α_S
- **Constrains on PDFs**
- SM phase diagram
- BSM searches in production and decay



Why precision predictions for top-quark pair production? → Applications


- top quark mass and production cross-sections
→ precision test of SM
- measurements of α_S
- Constrains on PDFs
- **SM phase diagram**
- BSM searches in production and decay






Why precision predictions for top-quark pair production? → Applications

- top quark mass and production cross-sections → precision test of SM
- measurements of α_S
- Constrains on PDFs
- SM phase diagram
- **BSM searches in production and decay**


New physics in top quark production



... and decay



Kinematics and polarization



External Momenta

$$p_1^2 = p_2^2 = 0$$

$$p_3^2 = p_4^2 = m^2$$

Mandelstamm variables

$$s = (p_1 + p_2)^2$$

$$t = (p_1 - p_3)^2$$

$$u = (p_2 - p_3)^2$$

$$s + t + u = 2m^2$$

Polarization sum external
gluons (axial gauge)

$$\sum_{\text{pol}} \epsilon_{i\mu}^* \epsilon_{i\nu} =$$

$$-g_{\mu\nu} + \frac{n_{i\mu} p_{i\nu} + n_{i\nu} p_{i\mu}}{n_i \cdot p_i}$$

Equation of motion for
external (anti)quarks

$$(\not{p} - m) U = 0$$

$$(\not{p} + m) V = 0$$

IBP reduction

General two-loop integral:

$$\int \frac{d^d l_1}{(2\pi)^d} \frac{d^d l_2}{(2\pi)^d} \prod_i \frac{1}{D_i^{n_i}} \prod_j N_j^{n_j}$$

with $D_i = (\sum p + \sum l)^2 - m^2$ and $N_i = l \cdot p$

Basic Idea of Integration-By-Part (IBP) reduction:

$$\int \frac{d^d l_1}{(2\pi)^d} \frac{d^d l_2}{(2\pi)^d} \frac{\partial}{\partial q^\mu} q^\mu I(l_1, l_2, \{p_{\text{ext}}\}) = 0 \text{ with } q = l_1, l_2, \{p_{\text{ext}}\}$$

- Relations between different integrals
⇒ Relate difficult integrals to easy ones
- Reduction to set of master integrals

Differential equations for master integrals

- Differentiation of master integrals with respect to m_s and x :
 $m_s \frac{d}{dm_s} I_i = \dots \quad x \frac{d}{dx} I_i = \dots$
- IBPs \rightarrow reduce the R.H.S again to masters
 \rightarrow coupled system of first order ODEs

$$m_s \frac{d}{dm_s} I_i = \sum c_k I_k$$

$$x \frac{d}{dx} I_i = \sum d_k I_k$$


- Boundary conditions \rightarrow Solution

Boundaries

- Analytic expansion around the high energy limes $m_s = \frac{M^2}{s} \rightarrow 0$
- Using Mellin-Barns representations and a lot of handwork to extract a series in $\epsilon, m_s s$ and x for each integral
- Expanding the differential equations also in ϵp and solve the algebraic system
- \rightarrow deep expansions in ms and x

Numerical evaluation of master integrals

- Using the differential equations to integrate numerically from the pre-calculated boundary conditions
- leaving the real numbers and integrate in a complex plane to grid points



The grid

Choice of points:

- $\beta = \sqrt{1 - 4m_s} = i/80$ for $i = 1, \dots, 79$
- 42 points for x : Gauss-Konrod points in available phase-space

