

Three photon production at the LHC: Amplitudes and Phenomenology

Milan - Bicocca seminar

Rene Poncelet

in collaboration with H. Chawdhry, M. Czakon and A. Mitov.

19th February 2020

Cavendish Laboratory




- Introduction
- Phenomenology
- Massless 5-point amplitudes at two-loop

Precision at the LHC

Standard Model Production Cross Section Measurements

Status:
November 2019

$\int \mathcal{L} dt$
[fb⁻¹]



Tremendous progress in NNLO QCD calculation in the past decade

State-of-the-art:

- All (Standard Model) $2 \rightarrow 2$ processes calculated
 - Phenomenology: SM precision measurements and parameter estimation, PDF determination, ...
- Valuable input for the LHC physics program!
- $2 \rightarrow 3$ is the natural step beyond. Many efforts on-going

Not quite comparable to the 'NLO revolution' yet, lack of automated:

1. Real radiation contributions → subtraction schemes
2. Two-loop matrix elements

Handling real radiation contribution in NNLO calculations cancellation of infra-red divergences

increasing number of available NNLO calculations with a variety of schemes

- **qT-slicing** [Catani,Grazzini, '07] , [Ferrera,Grazzini,Tramontano, '11], [Catani,Cieri,DeFlorian,Ferrera,Grazzini,'12], [Gehrmann,Grazzini,Kallweit,Maierhofer,Manteuffel,Rathlev,Torre,'14-'15], [Bonciani,Catani,Grazzini,Sargsyan,Torre,'14-'15], [Grazzini "MATRIX" '17-'19]
- **N-jettiness slicing** [Gaunt,Stahlhofen,Tackmann,Walsh, '15], [Boughezal,Focke,Giele,Liu,Petriello,'15-'16] , [Boughezal,Campell,Ellis,Focke,Giele,Liu,Petriello,'15], [Campell,Ellis,Williams,'16]
- **Antenna subtraction** [Gehrmann, GehrmannDeRidder,Glover,Heinrich,'05-'08] , [Weinzierl,'08,'09], [Currie,Gehrmann,GehrmannDeRidder,Glover,Pires,'13-'17], [Bernreuther,Bogner,Dekkers,'11,'14], [Abelof,(Dekkers),GehrmannDeRidder,'11-'15], [Abelof,GehrmannDeRidder,Maierhofer,Pozzorini,'14], [Chen,Gehrmann,Glover,Jaquier,'15]
- **Colorful subtraction** [DelDuca,Somogyi,Troscanyi,'05-'13], [DelDuca,Duhr,Somogyi,Tramontano,Troscanyi,'15]
- **Sector-improved residue subtraction (STRIPPER)** [Czakon,'10,'11] , [Czakon,Fiedler,Mitov,'13,'15], [Czakon,Heymes,'14] [Czakon,Fiedler,Heymes,Mitov,'16,'17], [Bughezal,Caola,Melnikov,Petriello,Schulze,'13,'14], [Bughezal,Melnikov,Petriello,'11], [Caola,Czernecki,Liang,Melnikov,Szafron,'14], [Bruchseifer,Caola,Melnikov,'13-'14], [Caola,Melnikov, Röntsch,'17-'19]
- **Projection-to-Born** [Cacciari et al '15], [Dreyer,Karlberg '18], **Geometric** [Herzog '18], **Unsubtraction** [Aguilera-Verdugo et al '19], . . .

Subtraction beyond $2 \rightarrow 2$ with the STRIPPER c++ framework

- Jet-production at NNLO QCD [Czakon,van Hameren,Mitov,Poncelet'19]: full set of subtraction terms in action
- Fully automated generation of subtraction terms
- Straight-forward user interface:
 - Generation of required contributions
 - Combination of equivalent contributions \rightarrow minimize computational setup
 - Automated interfaces to OpenLoops2 [Buccioni et al. '19] and Recola2 [Denner et al. '16-17], including available correlators

The framework is ready for the future

Introduction: Why $pp \rightarrow \gamma\gamma$?



- Potentially simplest $2 \rightarrow 3$ process:
 - Simple IR-structure
 - Allows for tool development
 - First experience with the two-loop matrix elements
 - Two-loop amplitudes: fewer diagrams and channels than jets

→ Proof-of-principle calculation

- Phenomenologically interesting:
 - LHC measurements show significant deviations from standard NLO predictions
 - Large NNLO/NLO K-factors have been observed in photon pair production
[Catani et al 11],[Campbell et al 16]

Introduction: Photon pair production





- Giant K-factor $\sigma^{\text{NNLO}}/\sigma^{\text{NLO}} \approx 2$
- Significant impact on shape of differential distribution
- Good agreement between NNLO QCD and data (CMS 7 TeV)



[Campbell, Ellis, Li, Williams 16]

Introduction: Three isolated photons

Detailed differential measurements by ATLAS [1712.07291 ATLAS]



NNLO QCD corrections to three-photon production at the LHC

[Chawdhry, Czakon, Mitov, Poncelet '19]


Setup ATLAS [1712.07291]:

- $E_T (= p_T)$ cut for the three photons: $E_{T,\gamma_1} > 27$ GeV, $E_{T,\gamma_2} > 22$ GeV, $E_{T,\gamma_3} > 15$ GeV
- Rapidity: All photons have $|\eta_\gamma| < 2.37$ (+exclusion of $1.37 < |\eta_\gamma| < 1.56$)
- Separation of photons: The angular distance between each two photons ΔR is required to be > 0.45
- Invariant mass: $m_{\gamma\gamma\gamma} > 50$ GeV
- Photon isolation: Using the Frixione [Frixione '98] isolation as indicated for the MadGraph@NLO setup. This means $R_0 = 0.4$, $E_T^{iso} > 10$ GeV and $\chi(R) = (1 - \cos(\Delta R))/(1 - \cos(\Delta R_0))$.
- PDF set: *NNPDF31_nnlo_as_0118*
- Scales:


$$\mu_0 = m_{T,\gamma\gamma\gamma} = \sqrt{p_\gamma^2 + (p_{\gamma,T})^2} \quad \text{with} \quad p_\gamma = \sum_{i=1}^3 p_{\gamma_i}$$

$$\mu_0 = H_T/4 = \frac{1}{4} \sum p_{\gamma_i,T}$$

3 Photons @ LHC: Fiducial cross section





- Huge K-factors:
 $\sigma^{\text{NLO}}/\sigma^{\text{LO}} \approx 2 - 3$
 $\sigma^{\text{NNLO}}/\sigma^{\text{NLO}} \approx 1.5$
- Significant improvement in description of data
- Scale dependence dominated by $gg/q\bar{q}'$ channels



3 Photons @ LHC: Differential distributions


- Not only normalization \rightarrow significant effects on the shape
- Remarkable agreement of measurement with NNLO QCD



3 Photons @ LHC: Perturbative convergence



$$pp \rightarrow \gamma\gamma$$

- Similar large K-factors in $\gamma\gamma$
[Catani, Cieri, de Florian, Ferrera, Grazzini 11]
[Campbell, Ellis, Li, Williams 16]
- $gg \rightarrow \gamma\gamma$ 1-loop box
 $\sim +10\%$ cross section
 +sizeable NLO corrections ($\Delta\sigma_{gg, n_f}^{N3LO}$)




$$pp \rightarrow \gamma\gamma\gamma$$

- gg channel contributes through $gg \rightarrow \gamma\gamma\gamma q\bar{q}$ @ NNLO QCD
- $gg \rightarrow \gamma\gamma\gamma$ 1-loop box does vanish (Furry's Theorem)



3 Photons @ LHC: Perturbative convergence

- $gg \rightarrow \gamma\gamma\gamma$ 1-loop box vanishes (Furry's Theorem)
 - But $gg \rightarrow \gamma\gamma\gamma g$ does not, and is separately finite N³LO contribution
- negligible!



$$pp \rightarrow \gamma\gamma\gamma$$

- NNLO QCD corrections are vital for comparisons of data with SM
 - Improved normalization
 - Very good agreement of differential cross section between NNLO QCD and ATLAS data
- Huge K-factors $\sigma^{\text{NNLO}}/\sigma^{\text{NLO}} \approx 1.5$
 - Very similar to Photon pair production
 - but without gg -box contribution
- Scale dependence driven by 'LO' contributions
 - $H_T/4$ dynamical scale choice (similar to the scale used in top-pair production)

$$pp \rightarrow \gamma\gamma\gamma$$

- NNLO QCD corrections are vital for comparisons of data with SM
 - Improved normalization
 - Very good agreement of differential cross section between NNLO QCD and ATLAS data
- Huge K-factors $\sigma^{\text{NNLO}}/\sigma^{\text{NLO}} \approx 1.5$
 - Very similar to Photon pair production
 - but without gg -box contribution
- Scale dependence driven by 'LO' contributions
 - $H_T/4$ dynamical scale choice (similar to the scale used in top-pair production)

What is about the other $2 \rightarrow 3$ processes?

Main bottle-neck are the two-loop amplitudes!

Five-point amplitudes in the IBP approach

State-of-art:

→ no general algorithm for numerical evaluation (like cut-based one-loop amps)

→ case-by-case study

- Massless case:
 - Masters known
 - planar: numerical implementation [Gehrmann et al '18]
 - non-planar: differential equations [Chicherin et al'19]
 - Reductions:
 - planar: known analytically [Chawdhry et al '18]
 - non-planar: work in progress ([Guan, Liu, Ma '19]?)
- Massless 5-point amplitudes:
 - $pp \rightarrow \gamma\gamma\gamma$ (N_c^3 contribution done)
 - $pp \rightarrow \gamma\gamma j$
 - $pp \rightarrow \gamma jj$
 - $pp \rightarrow jjj$
(planar, euclidean region only [Abreu,Dormans,Febres Cordero,Ita,Page,Sotnikov '19])
- Massive particles?
more scales \Leftrightarrow more difficult (a lot of unknowns: Reductions, Masters)

The traditional approach to amplitudes

Feynman diagrams



Tensor reduction \rightarrow scalar integrals



IBP reduction of scalar integrals to masters



Expression of masters in terms of function basis (Polylogarithms, etc.)



Evaluable (Semi-) Analytic expression of the matrix element

The traditional approach to amplitudes

Feynman diagrams



Tensor reduction \rightarrow scalar integrals



IBP reduction of scalar integrals to masters



Expression of masters in terms of function basis (Polylogarithms, etc.)



Evaluable (Semi-) Analytic expression of the matrix element


5-point 2-loop: $q\bar{q} \rightarrow \gamma\gamma\gamma$ matrix element

- Diagram generation with DiaGen [Czakon, private code] \rightarrow 1200 diagrams
- Color and fermion-loop structure of the matrix element:

$$\begin{aligned} \sum_{\text{I.C.}} 2\text{Re} \langle \mathcal{M}^{(0)} | \mathcal{M}^{(2)} \rangle = & \mathcal{M}^{(\text{lc},1)} C_F^2 C_A + \mathcal{M}^{(\text{lc},2)} C_F C_A^2 + \mathcal{M}^{(\text{f})} C_A C_F \\ & + \mathcal{M}^{(\text{np})} (N_c - 1/N_c) \end{aligned}$$

- Color decomposition in the leading colour approximation

$$\sum_{\text{I.C.}} 2\text{Re} \langle \mathcal{M}^{(0)} | \mathcal{M}^{(2)} \rangle \approx N_c^3 \left(\mathcal{M}^{(\text{lc},1)} + \mathcal{M}^{(\text{lc},2)} \right)$$



- neglecting $\mathcal{M}^{(\text{f})}$ contribution ($\sim n_f N_c^2$), contains non-planar contribution

5-point 2-loop: Scalar integrals

- All $k_1 \cdot k_2$ and $k_i \cdot p_j$ expressed through inverse propagators
- 11-propagator integral:

$$B[\vec{a}] = \left\{ k_1^2, k_2^2, (k_1 + p_1)^2, (k_1 + p_1 + p_2)^2, \right. \\ (k_2 - p_3)^2, (k_2 - k_1 - p_3)^2, \\ (k_2 - k_1 - p_1 - p_2 + p_4)^2, (k_2 + p_4)^2, \\ \left. (k_2 + p_1 + p_2)^2, (k_2 + p_1)^2, (k_1 + p_3)^2 \right\}$$

$$C[\vec{a}] = \left\{ k_1^2, k_2^2, (k_1 + p_1 + p_2)^2, (k_1 - k_2)^2, \right. \\ (k_2 + p_1)^2, (k_2 + p_1 + p_2)^2, (k_2 - p_3)^2, \\ (k_1 + p_1 + p_2 - p_3)^2, (k_1 + p_1 + p_2 - p_3 - p_4)^2, \\ \left. (k_2 - p_3 - p_4)^2, (k_1 + p_1)^2 \right\}$$

$$\Rightarrow \mathcal{M}^{(C)} = \sum c_i(\{s_{ij}\}, \epsilon) I(\vec{a}_i) \quad I \text{ either B or C}$$

The traditional approach to amplitudes

Feynman diagrams



Tensor reduction \rightarrow scalar integrals



IBP reduction of scalar integrals to masters



Expression of masters in terms of function basis (Polylogarithms, etc.)



Evaluable (Semi-) Analytic expression of the matrix element

Massless 5-point 2-loop: IBP identities and reduction

Topologies for massless 5-point amplitudes


$$B_1 = B [1, 1, 1, 1, 1, 1, 1, 1, 0, 0, 0]$$

$$B_2 = B [1, 1, 1, 1, 0, 1, 1, 1, 0, 0, 1]$$

$$C_1 = C [1, 1, 1, 1, 1, 1, 0, 1, 1, 0, 0]$$

$$C_2 = C [1, 1, 1, 1, 1, 0, 0, 1, 1, 0, 1]$$

$$C_3 = C [1, 0, 1, 1, 1, 1, 0, 1, 0, 0, 1]$$



- Reduction of planar topologies up to numerator power -5 available:

[Chawdhry, Lim, Mitov '18]

- Memory and CPU intensive venture
- 'divide and conquer': solve IBPs for one master at a time \rightarrow easy to parallelize and reduced memory consumption
- Non-planar topologies: work ongoing, but is constraint by available CPU hours, recent developments [Guan, Liu, Ma '19]

The traditional approach to amplitudes

Feynman diagrams



Tensor reduction \rightarrow scalar integrals



IBP reduction of scalar integrals to masters



Expression of masters in terms of function basis (Polylogarithms, etc.)



Evaluable (Semi-) Analytic expression of the matrix element

5-point 2-loop: Master Integrals

Master integrals expressed through planar 'pentagon-function'-basis

[Gehrmann,Henn,Presti '18]

- DEQ for masters $\vec{I} = \sum_k \epsilon^k \vec{I}^{(k)}$ in ϵ -form: $d\vec{I}^{(k+1)}(s_{ij}) = d\vec{A}(s_{ij})\vec{I}^{(k)}(s_{ij})$
- $\vec{I}^{(k)}$ can be expressed through iterated integrals
- Independent functions:

Weight	Functions
1	1
2	1
3	4
4	11

Note: Independent in the sense that $\ln(s_{12}) \leftrightarrow \ln(s_{23})$


- Different boundary constants for different physical regions

5-point 2-loop: Crossings

Large set of functions (~ 1800 @ Weight 4, including lower weight products)
due to momenta permutations

The difference to an euclidean amplitude (particularly in sub topologies).

Example:



5-point 2-loop: The amplitude in terms functions

$$\mathcal{M}^{(C)} = \sum_{k=-4}^0 \sum_l c_l^k(\{s_{ij}\}) f_l^k \epsilon^k$$

- Building amplitude together \rightarrow computationally intensive
- Large cancellation between in the rational coefficient $\mathcal{O}(1GB) \rightarrow \mathcal{O}(1MB)$ after simplifying
- Usage of rational reconstruction software FiniteFlow [Peraro '19] to sum up coefficients $c_l^k(\{s_{ij}\})$ (but under usage of the analytical reduction result)
- Cancellation of UV and IR poles checked analytically

5-point 2-loop: Evaluation

- Rational c++ implementation of coefficients with the help of CLN library to avoid loss numerical precision
- Usage of 'pentagon-function' implementation by [Gehrmann,Henn,Presti '18]
- 10 to 50 minutes per phase space point, 30k points evaluated (unweighted Born PS points)
- Coefficients relatively fast (~ 1 min), the functions take most of the time (numerical integration)
- Stability checks by changing integration precision
- Additional checks with interpolation software **GPTree** [Kasabov '19] to detect numerical instabilities

5-point 2-loop: Beyond $pp \rightarrow \gamma\gamma|N_c^3$

- Same technology applicable to the non-planar contributions (as soon as reductions are available, Master representation?)
- The other 5-point amplitudes:
 - $pp \rightarrow \gamma\gamma j$
 - $pp \rightarrow \gamma jj$
 - $pp \rightarrow jjj$

are feasible as well (at least in the leading colour limit)


- Combination with real radiation?
⇒ NNLO QCD within the STRIPPER framework
 - Challenges: more complicated radiation pattern due to final state colored particles, growing number of partonic contributions, stability and convergence?

The Future

- LHC-HL \Rightarrow more and more statistics which allow for:
 - Unprecedented precision in SM measurements
 - More exclusive selections
 - Rare processes measured qualitatively and quantitatively
- Precision predictions for multi-leg processes become more and more important!

Perturbative QCD is the backbone of theory predictions at the LHC

LHC/ High-Luminosity LHC timeline




Precision at the LHC

Standard Model Production Cross Section Measurements

Status:
November 2019

$\int \mathcal{L} dt$
[fb⁻¹]



2 \rightarrow 3 Phenomenology

- First process: $pp \rightarrow \gamma\gamma\gamma$
- Shows importance of NNLO QCD beyond $2 \rightarrow 2$
- Will certainly become more and more important with more experimental statistics



Advances for 5-point amplitudes:

- Application of IBP reductions for $pp \rightarrow \gamma\gamma\gamma$
- Finite remainder constructed and ready for use
- Certainly not the end of the story, many more amplitudes feasible with same techniques (5 partons, 4 partons + photon, 3 partons + 2 photons)

Backup



3 Photons @ LHC: Differential distributions

- Not only normalization \rightarrow significant effects on the shape
- Remarkable agreement of measurement with NNLO QCD





3 Photons @ LHC: Differential distributions

- Not only normalization \rightarrow significant effects on the shape
- Remarkable agreement of measurement with NNLO QCD



3 Photons @ LHC: Differential distributions

- Not only normalization \rightarrow significant effects on the shape
- Remarkable agreement of measurement with NNLO QCD




Refined formulation of the sector-improved residue subtraction

[Czakon '10 '11][Czakon,Heymes '14][Czakon,van Hameren,Mitov,Poncellet '19]

- New phase space parametrization:
 - Starts from Born kinematics \rightarrow additional radiation accommodated by rescaling and boosts
 - Generates minimal set of subtraction kinematics in each sector
 - Only one (!) double unresolved kinematic (= Born kinematic)
- Minimal set of sectors
- New 4-dimensional formulation:
 - New method to determine necessary counter terms
 - Numerical pole cancellation for each Born phase space point

Sector decomposition:



First complete NNLO QCD calculation for inclusive jet production

[Czakon,van Hameren,Mitov,Poncelet]

Many publications and studies by the NNLOJET collaboration:


[Currie, Gehrmann-De Ridder, Gehrmann, Glover, Huss, Pires '16-19]

- Antenna subtraction formalism
- Leading colour approximation for channels with quarks (expected to be a good approximation)
- Extensive analysis of renormalization scale setting and dependence:
 - Cancellation between different n-jet samples!
 - Distinguish 'jet'- and 'event'-type scales:
 - Inclusive jet observables: $\mu = p_T$ for each jet
- Very good description of LHC data for various observables: inclusive jets, various di-jet observables.

Technically very challenging process.
Contains the full set of NNLO IR singularities!

Stripper: Single-inclusive jet cross sections

- First full NNLO QCD calculation at 13 TeV
- Quite slow convergence: 350k CPU hours \rightarrow optimization potential!
- Comparison to NNLOJET: sub-leading colour effects within MC errors, thus indeed small
- K-factors public



Stripper: Single-inclusive jet cross sections

- First full NNLO QCD calculation at 13 TeV
- Quite slow convergence: 350k CPU hours \rightarrow optimization potential!
- Comparison to NNLOJET: sub-leading colour effects within MC errors, thus indeed small
- K-factors public

