

# Techniques and phenomenology of cutting-edge higher-order calculations for LHC processes

---

Rene Poncelet



THE HENRYK NIEWODNICZAŃSKI  
INSTITUTE OF NUCLEAR PHYSICS  
POLISH ACADEMY OF SCIENCES

Presented research received  
funding from:

LEVERHULME  
TRUST




# Outline

---


- Introduction
- Sector-improved residue subtraction
- Two-loop five-point amplitudes
- Pheno @ LHC:
  - Three-jet production through NNLO QCD
  - HighTEA
- Summary and Outlook

# What are the fundamental building blocks of matter?


Scattering experiments



Credit: CERN




Theory/Model



Credit: Jack Lindon, CERN

# SM measurements at the LHC



New physics around the corner?

Precise measurements  
 <->  
 Precise theory

- improved SM understanding
- search for indirect NP signals

# Precision predictions

---

**Fixed order  
perturbation theory**

Resummation

Parton-showers


Precision theory predictions

Parametric input:  
PDFs, couplings ( $\alpha_s$ ), ...

Soft physics:  
MPI, colour reconnection,  
...

Fragmentation/hadronisation

# Perturbative QCD



Hadronic cross section:

$$\sigma_{h_1 h_2 \rightarrow X} = \sum_{ij} \int_0^1 \int_0^1 dx_1 dx_2 \underbrace{\phi_{i,h_1}(x_1, \mu_F^2)}_{\text{PDF}} \underbrace{\phi_{j/h_2}(x_2, \mu_F^2)}_{\text{PDF}} \underbrace{\hat{\sigma}_{ij \rightarrow X}(\alpha_s(\mu_R^2), \mu_R^2, \mu_F^2)}_{\text{Partonic cross section}}$$

Parton distribution functions:  $\delta \sim 1\text{-}3\%$

Perturbative expansion of partonic cross section:

$$\hat{\sigma}_{ab \rightarrow X} = \hat{\sigma}_{ab \rightarrow X}^{(0)} + \hat{\sigma}_{ab \rightarrow X}^{(1)} + \hat{\sigma}_{ab \rightarrow X}^{(2)} + \mathcal{O}(\alpha_s^3)$$

Typical uncertainties from scale variations:  $\delta \text{LO} \sim \mathcal{O}(\sim 100\%)$   $\delta \text{NLO} \sim \mathcal{O}(\sim 10\%)$   $\delta \text{NNLO} (\sim 1\%)$   
 (estimate for corrections from missing higher orders)

# Next-to-leading order case

$$\hat{\sigma}_{ab}^{(1)} = \hat{\sigma}_{ab}^R + \hat{\sigma}_{ab}^V + \hat{\sigma}_{ab}^C$$




## KLN theorem

sum is finite for sufficiently inclusive observables and regularization scheme independent

Each term separately infrared (IR) divergent:


Real corrections:



$$\hat{\sigma}_{ab}^R = \frac{1}{2\hat{s}} \int d\Phi_{n+1} \langle \mathcal{M}_{n+1}^{(0)} | \mathcal{M}_{n+1}^{(0)} \rangle F_{n+1}$$

Phasespace integration over unresolved configurations

Virtual corrections:



$$\hat{\sigma}_{ab}^V = \frac{1}{2\hat{s}} \int d\Phi_n 2\text{Re} \langle \mathcal{M}_n^{(0)} | \mathcal{M}_n^{(1)} \rangle F_n$$

Integration over loop-momentum

Collinear factorization:  $\hat{\sigma}_{ab}^C = (\text{single convolution}) F_n$

# Slicing and Subtraction

Central idea: Divergences arise from infrared (IR, soft/collinear) limits → Factorization!

## Slicing

$$\hat{\sigma}_{ab}^R = \frac{1}{2\hat{s}} \int_{\delta(\Phi) \geq \delta_c} d\Phi_{n+1} \langle \mathcal{M}_{n+1}^{(0)} | \mathcal{M}_{n+1}^{(0)} \rangle F_{n+1} + \frac{1}{2\hat{s}} \int_{\delta(\Phi) < \delta_c} d\Phi_{n+1} \langle \mathcal{M}_{n+1}^{(0)} | \mathcal{M}_{n+1}^{(0)} \rangle F_{n+1}$$

$$\approx \frac{1}{2\hat{s}} \int_{\delta(\Phi) \geq \delta_c} d\Phi_{n+1} \langle \mathcal{M}_{n+1}^{(0)} | \mathcal{M}_{n+1}^{(0)} \rangle F_{n+1} + \frac{1}{2\hat{s}} \int d\Phi_n \tilde{M}(\delta_c) F_n + \mathcal{O}(\delta_c)$$

- Conceptually simple
- Recycling of lower computations
- Non-local cancellations/power-corrections  
→ computationally expensive

## Subtraction

$$\hat{\sigma}_{ab}^R = \frac{1}{2\hat{s}} \int \left( d\Phi_{n+1} \langle \mathcal{M}_{n+1}^{(0)} | \mathcal{M}_{n+1}^{(0)} \rangle F_{n+1} - d\tilde{\Phi}_{n+1} \mathcal{S}F_n \right) + \frac{1}{2\hat{s}} \int d\tilde{\Phi}_{n+1} \mathcal{S}F_n$$

$$\frac{1}{2\hat{s}} \int d\tilde{\Phi}_{n+1} \mathcal{S}F_n = \frac{1}{2\hat{s}} \int \underline{d\Phi_n d\Phi_1} \mathcal{S}F_n$$

- Conceptually more difficult
- Local subtraction → efficient
- Better numerical stability

Phasespace factorization  
→ momentum mappings

... +  $\hat{\sigma}_{ab}^V$  = finite



# Slicing and Subtraction

---

Central idea: Divergences arise from infrared (IR, soft/collinear) limits → Factorization!

## Slicing

qT-slicing [[Catain'07](#)],  
N-jettiness slicing [[Gaunt'15/Boughezal'15](#)]

- Conceptually simple
- Recycling of lower computations
- Non-local cancellations/power-corrections  
→ computationally expensive


## Subtraction

Antenna [[Gehrmann'05-'08](#)], Colorful [[DelDuca'05-'15](#)],  
Projection [[Cacciari'15](#)], Geometric [[Herzog'18](#)],  
Unsubtraction [[Aguilera-Verdugo'19](#)],  
Nested collinear [[Caola'17](#)],  
[Sector-improved residue subtraction](#) [[Czakon'10-'14'19](#)]

- Conceptually more difficult
- Local subtraction → efficient
- Better numerical stability


# Partonic cross section beyond NLO

$$\hat{\sigma}_{ab}^{(2)} = \hat{\sigma}_{ab}^{VV} + \hat{\sigma}_{ab}^{RV} + \hat{\sigma}_{ab}^{RR} + \hat{\sigma}_{ab}^{C2} + \hat{\sigma}_{ab}^{C1}$$




Real-Real

$$\hat{\sigma}_{ab}^{RR} = \frac{1}{2\hat{s}} \int d\Phi_{n+2} \langle \mathcal{M}_{n+2}^{(0)} | \mathcal{M}_{n+2}^{(0)} \rangle F_{n+2}$$



Real-Virtual

$$\hat{\sigma}_{ab}^{RV} = \frac{1}{2\hat{s}} \int d\Phi_{n+1} 2\text{Re} \langle \mathcal{M}_{n+1}^{(0)} | \mathcal{M}_{n+1}^{(1)} \rangle F_{n+1}$$




Virtual-Virtual

$$\hat{\sigma}_{ab}^{VV} = \frac{1}{2\hat{s}} \int d\Phi_n \left( 2\text{Re} \langle \mathcal{M}_n^{(0)} | \mathcal{M}_n^{(2)} \rangle + \langle \mathcal{M}_n^{(1)} | \mathcal{M}_n^{(1)} \rangle \right) F_n$$

$$\hat{\sigma}_{ab}^{C2} = (\text{double convolution}) F_n \quad \hat{\sigma}_{ab}^{C1} = (\text{single convolution}) F_{n+1}$$

## Sector-improved residue subtraction

---



# Sector decomposition I


Considering working in CDR:

→ Virtuals are usually done in this regularization:  $\hat{\sigma}_{ab}^{VV} = \sum_{i=-4}^0 c_i \epsilon^i + \mathcal{O}(\epsilon)$

→ Can we write the real radiation as such expansion?

→ Difficult integrals, analytical impractical (except very simple observables)!

→ Numerics not possible, integrals are divergent →  $\epsilon$ -poles!



How to extract these poles? → Sector decomposition!

**Divide and conquer** the phase space:

$$1 = \sum_{i,j} \left[ \sum_k \mathcal{S}_{ij,k} + \sum_{k,l} \mathcal{S}_{i,k;j,l} \right] \longrightarrow \hat{\sigma}_{ab}^{\text{RR}} = \frac{1}{2\hat{s}} \int d\Phi_{n+2} \sum_{i,j} \left[ \sum_k \mathcal{S}_{ij,k} + \sum_{k,l} \mathcal{S}_{i,k;j,l} \right] \langle \mathcal{M}_{n+2}^{(0)} | \mathcal{M}_{n+2}^{(0)} \rangle_{\text{F}_{n+2}}$$

# Sector decomposition II

Divide and conquer the phase space

- Each  $\mathcal{S}_{ij,k}/\mathcal{S}_{i,k;j,l}$  has simpler divergences:
  - Soft limits of partons I (and j)
  - Collinear w.r.t partons k (and l) of partons i (and j)
- Parametrization w.r.t. reference parton makes divergences explicit

$$\hat{\eta}_i = \frac{1}{2}(1 - \cos \theta_{ir}) \in [0, 1]$$

$$\hat{\xi}_i = \frac{u_i^0}{u_{\max}^0} \in [0, 1]$$

- Example: Splitting function

$$\sim \frac{1}{s_{r1}} P(z)$$

$$s_{r1} = (p_r + p_1)^2 = 2p_r^0 u_{\max}^1 \xi_1 \eta_1$$


$$\sim \frac{1}{\eta_1 \xi_1}$$

$$\mathcal{S}_{i,k} = \frac{1}{D_1 d_{i,k}} \quad D_1 = \sum_{ik} \frac{1}{d_{i,k}}$$

$$d_{i,k} = \frac{E_i}{\sqrt{s}} (1 - \cos \theta_{ik})$$

# Sector decomposition II – triple collinear factorization

Three particle invariant does not factorize trivially:



Double soft factorization:

$$\theta(u_1^0 - u_2^0) + \theta(u_2^0 - u_1^0)$$

$$(p_r + u_1 + u_2)^2 = 2p_r^0 (\xi_1 \eta_1 u_{\max}^1 + \xi_2 \eta_2 u_{\max}^2 + \xi_1 \xi_2 \frac{u_{\max}^1 u_{\max}^2}{p_r^0} \angle(u_1, u_2))$$

[Czakov'10, Caola'17]

# Sector decomposition III

Factorized singular limits in each sector:

$$\frac{1}{2\hat{s}} \int d\Phi_{n+2} \mathcal{S}_{kl,m} \langle \mathcal{M}_{n+2}^{(0)} | \mathcal{M}_{n+2}^{(0)} \rangle F_{n+2} = \sum_{\text{sub-sec.}} \int d\Phi_n \prod dx_i \underbrace{x_i^{-1-b_i\epsilon}}_{\text{singular}} d\tilde{\mu}(\{x_i\}) \underbrace{\prod x_i^{a_i+1} \langle \mathcal{M}_{n+2} | \mathcal{M}_{n+2} \rangle}_{\text{regular}} F_{n+2}$$

$x_i \in \{\eta_1, \xi_1, \eta_2, \xi_2\}$

Regularization of divergences:

$$x^{-1-b\epsilon} = \underbrace{\frac{-1}{b\epsilon}}_{\text{pole term}} + \underbrace{[x^{-1-b\epsilon}]_+}_{\text{reg. + sub.}} \quad \int_0^1 dx [x^{-1-b\epsilon}]_+ f(x) = \int_0^1 \frac{f(x) - f(0)}{x^{1+b\epsilon}}$$

# Finite NNLO cross section

$$\hat{\sigma}_{ab}^{RR} = \frac{1}{2\hat{s}} \int d\Phi_{n+2} \langle \mathcal{M}_{n+2}^{(0)} | \mathcal{M}_{n+2}^{(0)} \rangle F_{n+2}$$

$$\hat{\sigma}_{ab}^{C1} = (\text{single convolution}) F_{n+1}$$

$$\hat{\sigma}_{ab}^{RV} = \frac{1}{2\hat{s}} \int d\Phi_{n+1} 2\text{Re} \langle \mathcal{M}_{n+1}^{(0)} | \mathcal{M}_{n+1}^{(1)} \rangle F_{n+1}$$

$$\hat{\sigma}_{ab}^{C2} = (\text{double convolution}) F_n$$

$$\hat{\sigma}_{ab}^{VV} = \frac{1}{2\hat{s}} \int d\Phi_n \left( 2\text{Re} \langle \mathcal{M}_n^{(0)} | \mathcal{M}_n^{(2)} \rangle + \langle \mathcal{M}_n^{(1)} | \mathcal{M}_n^{(1)} \rangle \right) F_n$$



sector decomposition and master formula

$$x^{-1-b\epsilon} = \underbrace{\frac{-1}{b\epsilon}}_{\text{pole term}} + \underbrace{[x^{-1-b\epsilon}]_+}_{\text{reg. + sub.}}$$

$$(\sigma_F^{RR}, \sigma_{SU}^{RR}, \sigma_{DU}^{RR}) \quad (\sigma_F^{RV}, \sigma_{SU}^{RV}, \sigma_{DU}^{RV}) \quad (\sigma_F^{VV}, \sigma_{DU}^{VV}, \sigma_{FR}^{VV}) \quad (\sigma_{SU}^{C1}, \sigma_{DU}^{C1}) \quad (\sigma_{DU}^{C2}, \sigma_{FR}^{C2})$$



re-arrangement of terms  $\rightarrow$  4-dim. formulation [Czakon'14, Czakon'19]

$$\underline{(\sigma_F^{RR})} \quad \underline{(\sigma_F^{RV})} \quad \underline{(\sigma_F^{VV})} \quad \underline{(\sigma_{SU}^{RR}, \sigma_{SU}^{RV}, \sigma_{SU}^{C1})} \quad \underline{(\sigma_{DU}^{RR}, \sigma_{DU}^{RV}, \sigma_{DU}^{VV}, \sigma_{DU}^{C1}, \sigma_{DU}^{C2})} \quad \underline{(\sigma_{FR}^{RV}, \sigma_{FR}^{VV}, \sigma_{FR}^{C2})}$$

separately finite:  $\epsilon$  poles cancel



# C++ framework

---

- Formulation allows efficient algorithmic implementation → STRIPPER
- **High degree of automation:**
  - Partonic processes (taking into account all symmetries)
  - Sectors and subtraction terms
  - Interfaces to Matrix-element providers + O(100) hardcoded: AvH, OpenLoops, Recola, NJET, HardCoded
    - In practice: Only **two-loop matrix** elements required
- **Broad range of applications** through additional facilities:
  - Narrow-Width & Double-Pole Approximation
  - Fragmentation
  - Polarised intermediate massive bosons
  - (Partial) Unweighting → Event generation for **HighTEA**
  - Interfaces: FastNLO, FastJet

## Two-loop five-point amplitudes

---


Massless:

[Chawdry'19'20'21]  $(3A+2j, 2A+3j)$

[Abreu'20'21]  $(3A+2j, 5j)$

[Agarwal'21]  $(2A+3j)$

[Badger'21'23]  $(5j, gggAA, jjjA)$



1 external mass:

[Abreu'21]  $(W+4j)$


[Badger'21'22]  $(Hqqgg, W4q, Wajjj)$

[Hartanto'22]  $(W4q)$

# Overview

---


Old school approach:



Automated framework using finite fields  
to avoid expression swell based on  
FiniteFlow [Peraro'19]

# Projection to scalar integrals

Generate diagrams (contributing to leading-colour) with QGRAF



Factorizing decay:  $A_6^{(L)} = A_5^{(L)\mu} D_\mu P$

$$M_6^{2(L)} = \sum_{\text{spin}} A_6^{(0)*} A_6^{(L)} = M_5^{(L)\mu\nu} D_{\mu\nu} |P|^2$$

Projection on scalar functions (FORM+Mathematica):  
 → anti-commuting  $\gamma_5$  + Larin prescription

$$M_5^{(L)\mu\nu} = \sum_{i=1}^{16} a_i^{(L)} v_i^{\mu\nu}$$



$$a_i^{(L)} = a_i^{(L),\text{even}} + \text{tr}_5 a_i^{(L),\text{odd}}$$

$$a_i^{(L),p} = \sum_j c_{j,i}(\{p\}, \epsilon) \mathcal{I}(\{p\}, \epsilon)$$

# Integration-By-Parts reduction

$$a_i^{(L),p} = \sum_i c_{j,i}(\{p\}, \epsilon) \mathcal{I}(\{p\}, \epsilon) \quad \rightarrow \text{prohibitively large number of integrals}$$

$$\mathcal{I}_i(\{p\}, \epsilon) \equiv \mathcal{I}(\vec{n}_i, \{p\}, \epsilon) = \int \frac{d^d k_1}{(2\pi)^d} \frac{d^d k_2}{(2\pi)^d} \prod_{k=1}^{11} D_k^{-n_{i,k}}(\{p\}, \{k\})$$

Integration-By-Parts identities connect different integrals  $\rightarrow$  system of equations  
 $\rightarrow$  only a small number of independent "master" integrals

$$0 = \int \frac{d^d k_1}{(2\pi)^d} \frac{d^d k_2}{(2\pi)^d} l^\mu \frac{\partial}{\partial l^\mu} \prod_{k=1}^{11} D_k^{-n_{i,k}}(\{p\}, \{k\}) \quad \text{with } l \in \{p\} \cap \{k\}$$

LiteRed (+ Finite Fields)



$$a_i^{(L),p} = \sum_i d_{j,i}(\{p\}, \epsilon) \text{MI}(\{p\}, \epsilon)$$

# Master integrals & finite remainder

Differential Equations:  $d\vec{M}I = dA(\{p\}, \epsilon)\vec{M}I$

[Remiddi, 97]

[Gehrmann, Remiddi, 99]

[Henn, 13]

Canonical basis:  $d\vec{M}I = \epsilon d\tilde{A}(\{p\})\vec{M}I$

Simple iterative solution



$$MI_i = \sum_w \epsilon^w \tilde{M}I_i^w \quad \text{with} \quad \tilde{M}I_i^w = \sum_j c_{i,j} m_j$$

Chen-iterated integrals  
"Pentagon"-functions

[Chicherin, Sotnikov, 20]

[Chicherin, Sotnikov, Zoia, 21]




Putting everything together (and removing of IR poles):

$$f_i^{(L),p} = a_i^{(L),p} - \text{poles}$$

$$f_i^{(L),p} = \sum_j c_{i,j}(\{p\}) m_j + \mathcal{O}(\epsilon)$$

# Reconstruction of Amplitudes

[Badger'21]



## New optimizations

- Syzygy's to simplify IBPs
- Exploitation of Q-linear relations
- Denominator Ansatz
- On-the-fly partial fractioning


amplitude	helicity	original	stage 1	stage 2	stage 3	stage 4
$A_{34;q}^{(2),1}$	- + + - +	94/91	74/71	74/0	22/18	22/0
$A_{34;q}^{(2),1}$	- + - + +	93/89	90/86	90/0	24/14	18/0
$A_{34;q}^{(2),1/N_c^2}$	- + + - +	90/88	73/71	73/0	23/18	22/0
$A_{34;q}^{(2),1/N_c^2}$	- + - + +	90/86	86/82	86/0	24/14	19/0
$A_{34;l}^{(2),1/N_c}$	- + - + +	89/82	74/67	73/0	27/14	20/0
$A_{34;l}^{(2),1/N_c}$	- + + - +	85/81	61/58	60/0	27/18	20/0
$A_{34;q}^{(2),N_c^2}$	- + - + +	58/55	54/51	53/0	20/18	20/0

Massive reduction of complexity

# Overview

---

Old school approach:



Automated framework using finite fields  
to avoid expression swell based on  
FiniteFlow [Peraro'19]



# Three-jet production through NNLO QCD

---

# Multi-jet observables

## Test of pQCD and extraction of strong coupling constant

NLO theory unc. > experimental unc.

- **NNLO QCD needed for precise theory-data** comparisons  
→ Restricted to two-jet data [Currie'17+later][Czakon'19]
- **New NNLO QCD three-jet** → access to more observables
- Jet ratios



**Next-to-Next-to-Leading Order Study of Three-Jet Production at the LHC**  
Czakon, Mitov, Poncelet [2106.05331]

$$R^i(\mu_R, \mu_F, \text{PDF}, \alpha_{S,0}) = \frac{d\sigma_3^i(\mu_R, \mu_F, \text{PDF}, \alpha_{S,0})}{d\sigma_2^i(\mu_R, \mu_F, \text{PDF}, \alpha_{S,0})}$$


- Event shapes

**NNLO QCD corrections to event shapes at the LHC**

Alvarez, Cantero, Czakon, Llorente, Mitov, Poncelet [2301.01086]




# Encoding QCD dynamics in event shapes



Using (global) event information to separate different regimes of QCD event evolution:

- **Thrust & Thrust-Minor**  $T_{\perp} = \frac{\sum_i |\vec{p}_{T,i} \cdot \hat{n}_{\perp}|}{\sum_i |\vec{p}_{T,i}|}$ , and  $T_m = \frac{\sum_i |\vec{p}_{T,i} \times \hat{n}_{\perp}|}{\sum_i |\vec{p}_{T,i}|}$ .
- **Energy-energy correlators**

$$\frac{1}{\sigma_2} \frac{d\sigma}{d \cos \Delta\phi} = \frac{1}{\sigma_2} \sum_{ij} \int \frac{d\sigma_{x_{\perp,i} x_{\perp,j}}}{dx_{\perp,i} dx_{\perp,j} d \cos \Delta\phi_{ij}} \delta(\cos \Delta\phi - \cos \Delta\phi_{ij}) dx_{\perp,i} dx_{\perp,j} d \cos \Delta\phi_{ij},$$



Separation of energy scales:  $H_{T,2} = p_{T,1} + p_{T,2}$

Ratio to 2-jet:


$$R^i(\mu_R, \mu_F, \text{PDF}, \alpha_{S,0}) = \frac{d\sigma_3^i(\mu_R, \mu_F, \text{PDF}, \alpha_{S,0})}{d\sigma_2^i(\mu_R, \mu_F, \text{PDF}, \alpha_{S,0})}$$

Here: **use jets as input** → experimentally advantageous  
(better calibrated, smaller non-pert.)


# Transverse Thrust @ NNLO QCD

## NNLO QCD corrections to event shapes at the LHC

Alvarez, Cantero, Czakon, Llorente, Mitov, Poncelet [2301.01086]




ATLAS [2007.12600]



# The transverse energy-energy correlator

$$\frac{1}{\sigma_2} \frac{d\sigma}{d \cos \Delta\phi} = \frac{1}{\sigma_2} \sum_{ij} \int \frac{d\sigma_{x_{\perp,i}x_{\perp,j}}}{dx_{\perp,i}dx_{\perp,j}d \cos \Delta\phi_{ij}} \delta(\cos \Delta\phi - \cos \Delta\phi_{ij}) dx_{\perp,i} dx_{\perp,j} d \cos \Delta\phi_{ij},$$

- Insensitive to soft radiation through energy weighting  $x_{T,i} = E_{T,i} / \sum_j E_{T,j}$
- Event topology separation:
  - Central plateau contain isotropic events
  - To the right: self-correlations, collinear and in-plane splitting
  - To the left: back-to-back



[ATLAS 2301.09351]

**ATLAS**

Particle-level TEEC

$\sqrt{s} = 13 \text{ TeV}; 139 \text{ fb}^{-1}$

anti- $k_t$   $R = 0.4$

$p_T > 60 \text{ GeV}$

$|\eta| < 2.4$

$\mu_{R,F} = \hat{P}_T$


$\alpha_s(m_Z) = 0.1180$

NNPDF 3.0 (NNLO)

—•— Data  
 - - - LO  
 - . - . NLO  
 - - - NNLO

# Double differential TEEC

[ATLAS 2301.09351]



**ATLAS**

Particle-level TEEC

$\sqrt{s} = 13 \text{ TeV}; 139 \text{ fb}^{-1}$

anti- $k_t$   $R = 0.4$

$p_T > 60 \text{ GeV}$

$|\eta| < 2.4$

$\mu_{R,F} = \hat{p}_T$

$\alpha_s(m_Z) = 0.1180$

NNPDF 3.0 (NNLO)

— Data


--- LO

--- NLO


--- NNLO

# Running of $\alpha_S$

NLO QCD




NNLO QCD



# HighTEA

---





= ~100 MCPUh



How to make this more  
efficient/environment-friendly/  
accessible/faster?

high tea  
for your freshly brewed analysis

<https://www.precision.hep.phy.cam.ac.uk/hightea>

HighTEA: High energy Theory Event Analyser  
[2304.05993]

Michał Czakon,<sup>a</sup> Zahari Kassabov,<sup>b</sup> Alexander Mitov,<sup>c</sup> Rene Poncelet,<sup>c</sup> Andrei Popescu<sup>c</sup>

<sup>a</sup>*Institut für Theoretische Teilchenphysik und Kosmologie, RWTH Aachen University, D-52056 Aachen, Germany*

<sup>b</sup>*DAMTP, University of Cambridge, Wilberforce Road, Cambridge, CB3 0WA, United Kingdom*

<sup>c</sup>*Cavendish Laboratory, University of Cambridge, Cambridge CB3 0HE, United Kingdom*

*E-mail: [mczakon@physik.rwth-aachen.de](mailto:mczakon@physik.rwth-aachen.de), [zk261@cam.ac.uk](mailto:zk261@cam.ac.uk), [adm74@cam.ac.uk](mailto:adm74@cam.ac.uk), [poncelet@hep.phy.cam.ac.uk](mailto:poncelet@hep.phy.cam.ac.uk), [andrei.popescu@cantab.net](mailto:andrei.popescu@cantab.net)*

# Basic idea

---

## → Database of precomputed “Theory Events”

- **Equivalent to a full fledged computation**
- Currently this means partonic fixed order events
- Extensions to include showered/resummed/hadronized events is feasible
- (Partially) Unweighting to increase efficiency

Not so new idea:  
LHE [Alwall et al '06],  
Ntuple [BlackHat '08'13],

## → Analysis of the data through an user interface

- Easy-to-use
- Fast
  - Observables from basic 4-momenta
  - Free specification of bins
- Flexible:
  - Renormalization/Factorization Scale variation
  - PDF (member) variation
  - Specify phase space cuts

# Factorizations

---

Factorizing renormalization and factorization scale dependence:

$$w_s^{i,j} = w_{\text{PDF}}(\mu_F, x_1, x_2) w_{\alpha_s}(\mu_R) \left( \sum_{i,j} c_{i,j} \ln(\mu_R^2)^i \ln(\mu_F^2)^j \right)$$

PDF dependence:


$$w_{\text{PDF}}(\mu, x_1, x_2) = \sum_{ab \in \text{channel}} f_a(x_1, \mu) f_b(x_2, \mu)$$

$\alpha_s$  dependence:

$$w_{\alpha_s}(\mu) = (\alpha_s(\mu))^m$$


Allows **full control over scales and PDF**

# HighTEA interface




# Available Processes

Processes **currently** implemented in our STRIPPER framework through **NNLO QCD**



# The Vision


---



# Summary & Outlook

---

# The NNLO QCD revolution





# Take home message

---

- Subtraction at NNLO QCD gains maturity, challenges remain...
  - Efficiency!
  - Still difficult to run codes!
  - HighTEA possible solution?!
- Two-loop matrix elements for high multiplicity are the single most significant bottleneck for NNLO QCD calculations
  - 2 to 3 massless ME completed in full colour (~10 years of work of ~5 research groups)
  - 1 mass MEs next challenge...
- NNLO QCD is a staple for SM precision phenomenology
  - Challenge: matching to parton-shower!

# Backup

---

# Improved phase space generation

---

Phase space cut and differential observable introduce

*mis-binning*: mismatch between kinematics in subtraction terms

→ leads to increased variance of the integrand

→ slow Monte Carlo convergence

New phase space parametrization [Czakon'19]:

Minimization of # of different subtraction kinematics in each sector

# Improved phase space generation

New phase space parametrization:

Minimization of # of different subtraction kinematics in each sector


Mapping from  $n+2$  to  $n$  particle phase space:  $\{P, r_j, u_k\} \rightarrow \{\tilde{P}, \tilde{r}_j\}$

Requirements:

- Keep direction of reference  $r$  fixed
- Invertible for fixed  $u_i$ :  $\{\tilde{P}, \tilde{r}_j, u_k\} \rightarrow \{P, r_j, u_k\}$
- Preserve Born invariant mass:  $q^2 = \tilde{q}^2, \tilde{q} = \tilde{P} - \sum_{j=1}^{n_{fr}} \tilde{r}_j$

Main steps:

- Generate Born configuration
- Generate unresolved partons  $u_i$
- Rescale reference momentum  $r = x\tilde{r}$
- Boost non-reference momenta of the Born configuration



# Improved phase space generation

New phase space parametrization:

Minimization of # of different subtraction kinematics in each sector


Mapping from  $n+2$  to  $n$  particle phase space:  $\{P, r_j, u_k\} \rightarrow \{\tilde{P}, \tilde{r}_j\}$

Requirements:

- Keep direction of reference  $r$  fixed
- Invertible for fixed  $u_i$ :  $\{\tilde{P}, \tilde{r}_j, u_k\} \rightarrow \{P, r_j, u_k\}$
- Preserve Born invariant mass:  $q^2 = \tilde{q}^2, \tilde{q} = \tilde{P} - \sum_{j=1}^{n_{fr}} \tilde{r}_j$

Main steps:

- Generate Born configuration
- Generate unresolved partons  $u_i$
- Rescale reference momentum  $r = x\tilde{r}$
- Boost non-reference momenta of the Born configuration



# Improved phase space generation

New phase space parametrization:

Minimization of # of different subtraction kinematics in each sector


Mapping from  $n+2$  to  $n$  particle phase space:  $\{P, r_j, u_k\} \rightarrow \{\tilde{P}, \tilde{r}_j\}$

Requirements:

- Keep direction of reference  $r$  fixed
- Invertible for fixed  $u_i$ :  $\{\tilde{P}, \tilde{r}_j, u_k\} \rightarrow \{P, r_j, u_k\}$
- Preserve Born invariant mass:  $q^2 = \tilde{q}^2, \tilde{q} = \tilde{P} - \sum_{j=1}^{n_{fr}} \tilde{r}_j$

Main steps:

- Generate Born configuration
- Generate unresolved partons  $u_i$
- Rescale reference momentum  $r = x\tilde{r}$
- Boost non-reference momenta of the Born configuration



# Improved phase space generation

New phase space parametrization:

Minimization of # of different subtraction kinematics in each sector

Mapping from  $n+2$  to  $n$  particle phase space:  $\{P, r_j, u_k\} \rightarrow \{\tilde{P}, \tilde{r}_j\}$

Requirements:

- Keep direction of reference  $r$  fixed
- Invertible for fixed  $u_i$ :  $\{\tilde{P}, \tilde{r}_j, u_k\} \rightarrow \{P, r_j, u_k\}$
- Preserve Born invariant mass:  $q^2 = \tilde{q}^2, \tilde{q} = \tilde{P} - \sum_{j=1}^{n_{fr}} \tilde{r}_j$

Main steps:

- Generate Born configuration
- Generate unresolved partons  $u_i$
- Rescale reference momentum  $r = x\tilde{r}$
- Boost non-reference momenta of the Born configuration

